

$\Lambda(1405)N \rightarrow YN$ transition in nuclear medium for non-mesonic absorption of $\bar{K}$ in nucleus

T. Sekihara^a, D. Jido^b, Y. Kanada-En'yo^b

^aDepartment of Physics, Kyoto University, Kyoto 606-8502, Japan

^bYukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

Abstract

Non-mesonic transition of $\Lambda(1405)N \rightarrow YN$ is investigated as one of the essential processes for the non-mesonic absorption of $\bar{K}$ in nuclei. Using one-meson exchange model in the calculation of the transition, we find that the non-mesonic transition ratio $\Gamma_{\Lambda N}/\Gamma_{\Sigma N}$ depends strongly on the ratio of the $\Lambda(1405)$ (Λ^*) couplings to $\bar{K}N$ and $\pi\Sigma$. Especially a larger $\Lambda^*-\bar{K}N$ coupling leads to enhancement of the transition to ΛN . Using the chiral unitary model for the description of the Λ^* , we obtain $\Gamma_{\Lambda N}/\Gamma_{\Sigma N} \approx 1.2$ which is almost independent of the nucleon density, and find the total non-mesonic decay width of the Λ^* in uniform nuclear matter to be 22 MeV at the normal density.

Key words: Non-mesonic decay, kaonic nuclei; $\Lambda(1405)$ doorway; Chiral unitary approach

PACS: 21.85.+d, 14.20.Jn, 13.75.Jz, 13.30.Eg

1. Introduction

The study of the in-medium properties of anti-kaon ($\bar{K}$) has attracted continuous attention [1]. $\bar{K}$ has been theoretically expected to be bound by nuclear systems due to attractive strong interaction between $\bar{K}$ and nucleon N [2]. These $\bar{K}$ -nuclear bound systems assisted mainly by the strong interaction, which are so-called kaonic nuclei, have at least two important aspects; the kaonic nuclei are strongly interacting exotic many-body systems, and they provide favorable systems for the studies of the $\bar{K}$ properties at finite density. However, in spite of experimental efforts to search for the $\bar{K}$ -nuclear bound systems [3], there are no clear evidences observed yet.

In order to understand the $\bar{K}$ -nucleus systems, it is important to investigate theoretically the decay mechanism of the kaonic nuclei, or the absorption process of the $\bar{K}$ into nuclei. Especially the decay (absorption) ratios of various modes and explicit numbers of the decay (absorption) widths are key quantities for the $\bar{K}$ -nucleus systems. The decay of the kaonic nuclei in strong interactions can be categorized into two processes; one is the mesonic process such as $\bar{K}N \rightarrow \pi Y$, and the other is the non-mesonic process such as $\bar{K}NN \rightarrow YN$, where Y denotes hyperon (Λ or Σ). The non-mesonic decays have advantage in experimental observations, since signals from kaonic nuclei are readily distinguished from backgrounds and no extra mesons do not have to be detected. Therefore, systematic studies of the non-mesonic decay of the kaonic nuclei are desirable. Especially the ratios of the decay widths are interesting, since they will be insensitive to details of the production mechanism.

The absorption of the $\bar{K}$ in nuclei may take place dominantly through the $\Lambda(1405)$ (Λ^*) resonance, owing to the presence of the Λ^* just below the $\bar{K}N$ threshold. Namely the Λ^* can




Figure 1: Feynman diagrams for the $\Lambda^* N \rightarrow YN$ transition in the one-meson exchange model. The left two diagrams are for the ΛN final state and the right two are for the ΣN state.

be a doorway of the $\bar{K}$ absorptions in nuclei. The Λ^* doorway picture is more probable, in case that the Λ^* is a quasi-bound state of $\bar{K}N$ [4, 5], which has large $\bar{K}N$ components as almost real particles. Therefore the strong $\bar{K}N$ correlations are expected to be responsible for the Λ^* -induced decays of the $\bar{K}$ in nuclei. Motivated by the Λ^* doorway picture, we study the non-mesonic transition of $\Lambda^* N \rightarrow YN$ ($\Lambda^* p \rightarrow \Lambda p, \Sigma^0 p, \Sigma^+ n$ and $\Lambda^* n \rightarrow \Lambda n, \Sigma^0 n, \Sigma^- p$) in uniform nuclear matter with a one-meson exchange approach.

2. $\Lambda(1405)$ doorway process for the non-mesonic decay in kaonic nuclei

Now let us define the transition rate of the $\Lambda^* N \rightarrow YN$ process by the transition probability divided by time $\mathcal{T}$ as,

$$\gamma_{YN} \equiv \frac{1}{\mathcal{T}} \frac{1}{\mathcal{V}^2} \frac{1}{4} \sum_{\text{spin}} \int d\Phi_2 |S - 1|^2 = \frac{1}{\mathcal{V}} \frac{1}{4} \sum_{\text{spin}} \int d\Phi_2 |T_{YN}|^2 (2\pi)^4 \delta^4(P_{\Lambda^*} + P_{\text{in}} - P_N - P_Y), \quad (1)$$

with the S -matrix S for the transition process given by the transition amplitudes T_{YN} as $S = 1 - i(2\pi)^4 \delta^4(P_{\Lambda^*} + P_{\text{in}} - P_N - P_Y) T_{YN}$. $\mathcal{V}$ is volume of the system, and $d\Phi_2$ is the phase-space for the final state. The transition rate depends on the center-of-mass energy $E_{\text{c.m.}}$, equivalently the initial nucleon momentum. Summing up the $\Lambda^* N \rightarrow YN$ transition rate in terms of the initial nucleon states, we can estimate the non-mesonic decay width of the Λ^* in nuclear medium as a function of the nucleon density. The factor $1/\mathcal{V}$ in the last form of Eq. (1) is responsible for the fact that only one N exists in the initial state in the volume $\mathcal{V}$.

We evaluate the transition amplitudes T_{YN} with one-meson exchange diagrams shown in Fig. 1. The each diagram is composed of three parts: the s -wave $\Lambda^* MB$ coupling G_{MB} , the meson propagator including short-range correlation $\tilde{\Pi}(q^2)$ [7, 8], and the p -wave MBB coupling V_{MBB} , which is determined by the flavor SU(3) symmetry. The explicit form is given in Ref. [6].

Among the three parts for the transition, only the s -wave $\Lambda^* MB$ coupling constants G_{MB} are the model parameters, which are determined by the properties of the Λ^* . Therefore, in order to study the non-mesonic decay pattern in the Λ^* doorway, let us discuss the G_{MB} coupling dependence of the ratio of the transition rates, $\gamma_{\Lambda N}/\gamma_{\Sigma^0 N}$. With the conditions that $G_{\eta\Lambda} = 0$, the initial nucleon momentum $p_{\text{in}} = 0$, and the Λ^* mass $M_{\Lambda^*} = 1420$ MeV, we show the numerical result of the ratio of the transition rates, $\gamma_{\Lambda N}/\gamma_{\Sigma^0 N}$ as a function of $G_{\bar{K}N}/G_{\pi\Sigma}$ in Fig. 2, in which we assume $G_{\bar{K}N}/G_{\pi\Sigma}$ to be a real number. As seen in the figure, the ratio $\gamma_{\Lambda N}/\gamma_{\Sigma^0 N}$ has strong dependence on the coupling ratio $G_{\bar{K}N}/G_{\pi\Sigma}$. This is because the transition to ΛN is governed by the $\bar{K}$ exchange and the transition to ΣN is dominated by the π exchange due to the coupling V_{MBB} strength. This result suggests that larger $\Lambda^* \bar{K}N$ coupling leads to enhancement of the decay ratio to ΛN in the kaonic nuclei, although the Λ^* in vacuum cannot decay into final states including Λ .




Figure 2: Ratio of the transition rates of Λ^*N to ΛN and $\Sigma^0 N$, $\gamma_{\Lambda N}/\gamma_{\Sigma^0 N}$, as a function of $G_{\bar{K}N}/G_{\pi\Sigma}$. We fix the initial nucleon momentum $p_{\text{in}} = 0$ and the Λ^* mass $M_{\Lambda^*} = 1420$ MeV and assume $G_{\eta\Lambda} = 0$.

Now we calculate the non-mesonic decay of the Λ^* in uniform nuclear matter induced by the $\Lambda^*N \rightarrow YN$ transition, under the free Fermi gas approximation for nuclear matter. We evaluate the non-mesonic decay width Γ_{YN} by summing up the transition rate γ_{YN} for the nucleons:

$$\Gamma_{YN} \equiv \sum_{i=1}^{A_N} \gamma_{YN}(k_i) = \int_0^{k_{\text{FN}}} dk \frac{k^2}{\pi^2} \mathcal{V} \gamma_{YN}(k), \quad (2)$$

where $A_N = k_{\text{FN}}^3/(3\pi^2) \times \mathcal{V}$ ($N = p$ or n) is the numbers of the protons or neutrons in $\mathcal{V}$.

Furthermore we fix the ratios of the Λ^*MB coupling constants from the s -wave $\bar{K}N(I=0) \rightarrow MB$ (particle-basis) scattering amplitudes $t_{MB}(W)$ around the Λ^* resonance energy, as ratios of the $t_{MB}(W)$. Here W is the center-of-mass energy chosen to be the Λ^* resonance position on the real energy axis. For the description of the $\bar{K}N$ scattering amplitudes $t_{MB}(W)$, we use the chiral unitary approach (ChUA), which reproduces well the s -wave scattering amplitudes of $\bar{K}N \rightarrow MB$ in the coupled-channels method based on chiral dynamics with dynamically generated Λ^* [9]. We use the parameter set in Ref. [10] and take $W = M_{\Lambda^*} = 1420$ MeV, which is obtained by chiral unitary approach.

Fixing the ratios of the Λ^* coupling constants, G_{MB} , we plot the ratio of the non-mesonic decay widths of the in-medium $\bar{K}$ to ΛN and $\Sigma^0 N$ as a function of the proton (or neutron) density




Figure 3: (a) Non-mesonic decay width ratio of Λ^* in nuclei. (b) Non-mesonic decay width of Λ^* in nuclei. In both cases the Λ^* coupling constants are determined by the chiral unitary approach.

in Fig. 3(a). This figure shows that the ratio of the non-mesonic decay widths $\Gamma_{\Lambda N}/\Gamma_{\Sigma^0 N}$ is around 1.2 almost independently of the nucleon density. The density independence of the ratio of the decay widths is caused by sufficiently large phase-space in the final states.

We also obtain the absolute values of the non-mesonic decay widths of the Λ^* in nuclear matter. We use the coupling constants of the Λ^* to $\bar{K}N$, $\pi\Sigma$ and $\eta\Lambda$ obtained by the chiral unitary approach, which gives the in-vacuum mesonic decay as $\Gamma_{\Lambda^* \rightarrow \pi\Sigma} = 40$ MeV with $M_{\Lambda^*} = 1420$ MeV as observed in a K^- initiated channel [11, 12]. We show the non-mesonic decay widths in Fig. 3(b). The linear dependence of the decay widths is caused by the large phase-space in the final states. At the normal nuclear density ($\rho_B = 0.17 \text{ fm}^{-3}$), the total non-mesonic decay width is 22 MeV, which is almost half of the mesonic decay width of Λ^* (~ 40 MeV).

3. Conclusion

We have investigated the non-mesonic transition $\Lambda^* \bar{Y} \rightarrow YN$ in nuclear medium as one of the essential processes for the non-mesonic absorption of $\bar{K}$ in nucleus. Calculating the $\Lambda^* N \rightarrow YN$ transition rates in the one-meson exchange processes, we have found that the ratio of the $\Lambda^* N$ transition rates to ΛN and ΣN strongly depends on the ratio of the Λ^* couplings, $G_{\bar{K}N}/G_{\pi\Sigma}$. Especially, larger Λ^* couplings to $\bar{K}N$ lead to enhancement of the non-mesonic Λ^* decay with ΛN emission. Furthermore, describing the Λ^* properties by the chiral unitary approach, we have obtained the ratio of the non-mesonic decay widths to ΛN and $\Sigma^0 N$ as $\Gamma_{\Lambda N}/\Gamma_{\Sigma^0 N} = 1.2$ almost independently of the nucleon density. We have also estimated that the total non-mesonic decay width is 22 MeV at the saturation density. This study can be extend to the calculation of the absorption width of the $\bar{K}$ in nuclear medium [13].

References

- [1] C. J. Batty, E. Friedman and A. Gal, Phys. Rept. **287** (1997) 385; T. Waas and W. Weise, Nucl. Phys. **A625** (1997) 287; A. Ramos and E. Oset, Nucl. Phys. **A671** (2000) 481.
- [2] T. Kishimoto, Phys. Rev. Lett. **83** (1999) 4701; E. Friedman and A. Gal, Phys. Lett. B **459** (1999) 43; Y. Akaishi and T. Yamazaki, Phys. Rev. C **65** (2002) 044005; E. Friedman and A. Gal, Phys. Rept. **452** (2007) 89.
- [3] T. Kishimoto *et al.*, Prog. Theor. Phys. Suppl. **149** (2003) 264; M. Agnello *et al.* [FINUDA Collaboration], Phys. Rev. Lett. **94** (2005) 212303; T. Kishimoto *et al.*, Nucl. Phys. A **754** (2005) 383; T. Suzuki *et al.*, Nucl. Phys. A **754** (2005) 375.
- [4] R. H. Dalitz, T. C. Wong and G. Rajasekaran, Phys. Rev. **153** (1967) 1617.
- [5] T. Hyodo, D. Jido and A. Hosaka, Phys. Rev. C **78** (2008) 025203.
- [6] T. Sekihara, D. Jido and Y. Kanada-En'yo, Phys. Rev. C **79** (2009) 062201.
- [7] E. Oset and W. Weise, Nucl. Phys. A **319** (1979) 477.
- [8] D. Jido, E. Oset and J. E. Palomar, Nucl. Phys. A **694** (2001) 525.
- [9] N. Kaiser, P. B. Siegel and W. Weise, Nucl. Phys. A **594** (1995) 325; E. Oset and A. Ramos, Nucl. Phys. A **635** (1998) 99; E. Oset, A. Ramos and C. Bennhold, Phys. Lett. B **527** (2002) 99; D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meissner, Nucl. Phys. A **725** (2003) 181.
- [10] T. Sekihara, T. Hyodo and D. Jido, Phys. Lett. B **669** (2008) 133.
- [11] O. Braun *et al.*, Nucl. Phys. B **129** (1977) 1.
- [12] D. Jido, E. Oset and T. Sekihara, Eur. Phys. J. A **42** (2009) 257.
- [13] T. Sekihara, J. Yamagata-Sekihara, D. Jido and Y. Kanada-En'yo, in preparation.